

Daily Checklist for All Child Care Facilities

| Task | N/A | Completed | Date | Initials |
|---|-----|-----------|------|----------|
| Communication | | | | |
| Post signs at entry and exit regarding avoiding ill people, covering cough, | | | | |
| minimizing non-essential activities in the community, and frequent | | | | |
| handwashing | | | | |
| Stay in regular communication with all parents, children, staff, and | | | | |
| volunteers about steps everyone can take to protect themselves and | | | | |
| prevent further spread. | | | | |
| Provide staff and family means to contact facility for updates and concerns | | | | |
| Education and Preparation | | | | |
| Educate staff and families about: | | | | |
| 1. Signs and symptoms of COVID-19 | | | | |
| 2. Hand hygiene | | | | |
| 3. Reducing stigma | | | | |
| Review and update your emergency operations plan so that you can be | | | | |
| best prepared for cases of COVID-19. | | | | |
| Control Measures | | | | |
| Ask parents to check their children's temperature and ask staff to check | | | | |
| their own temperature daily before arrival at child care | | | | |
| If the temperature exceeds 100.4°F or if cough is present, advise | | | | |
| parents to isolate their child at home and ask staff members to isolate | | | | |
| at home. | | | | |
| Symptomatic staff and children should self isolate and not return | | | | |
| to the facility until 7 days after symptom onset OR 72 hours after | | | | |
| their fever is gone and initial symptoms have improved, whichever | | | | |
| is longer. | | | | |
| Establish daily procedures for identifying and sending home children | | | | |
| and staff who become sick (i.e., determine where child will wait for | | | | |
| parent, who will take child to meet parent). | | | | |
| Choose activities that allow for more physical space between children. | | | | |
| Increase frequency of <u>cleaning</u> , <u>sanitizing</u> , <u>and disinfecting</u> your child care | | | | |
| center. | | | | |
| Most common EPA-registered household disinfectants should be | | | | |
| effective. A list of products that are EPA-approved for use against the | | | | |
| virus that causes COVID-19 is available <u>here</u> . Products other than | | | | |
| bleach should be approved by licensor before use. | | | | |
| Establish and maintain hand hygiene stations stocked with either | | | | |
| soap/water/towels or alcohol-based hand sanitizer with a concentration of | | | | |
| at least 60% alcohol. These should be available outdoors as well. | | | | |



Checklist of ADDITIONAL Actions when a COVID Case has Occurred at a Child Care Facility

| Task | N/A | Completed | Date | Initials |
|---|-----|-----------|------|----------|
| Communication | | | | |
| Notify facility administration of COVID case. | | | | |
| Meet with key staff, including teachers, janitors and leadership, to | | | | |
| coordinate control measures. | | | | |
| Notify families and staff of a confirmed/potential COVID-19 | | | | |
| infection in the facility | | | | |
| Remember families should not be told personal identifiers | | | | |
| If possible, notify anyone at your child care setting that | | | | |
| the person infected with COVID-19 had close interactions | | | | |
| with while symptomatic. Make sure any close contacts are | | | | |
| being closely monitored for symptoms. | | | | |
| Infection Control Activities | | | | |
| Consider short-term closures as needed for cleaning | | | | |
| Clean, and disinfect the entire facility with EPA-registered | | | | |
| household disinfectants. Wait as long as practical before | | | | |
| beginning cleaning and disinfection of areas used by the | | | | |
| person infected with COVID-19 to minimize potential for | | | | |
| exposure to respiratory droplets. If possible, wait up to 24 | | | | |
| hours before beginning cleaning and disinfection. | | | | |
| If the individual infected with COVID-19 spent time in your child | | | | |
| care and had close contact with others while ill, you will need to | | | | |
| close for 14 days. Ensure staff and families understand ill people should remain | | | | |
| home until well and those with COVID-19-like symptoms should | | | | |
| self-isolate until 7 days after symptom onset OR 72 hours after | | | | |
| their fever is gone and initial symptoms have improved, whichever | | | | |
| is longer. | | | | |
| | | | | |
| Continued Service | | | | |
| Ensure continuity of meal programs when possible and consider | | | | |
| ways to distribute food to students. Consider options such as | | | | |
| "grab-and-go" bagged lunches or meal delivery. | | | | |